



« **Verbeterde doeltreffendheid deur innovasie** »



**VEGGIE  
OEMFF®**



## Naam:

**Veggies OEMFF® Starter, Veggies OEMFF® Grow, Veggies OEMFF® Fruit**

## Eienskappe (Wat):

'n Fyn, droë hoogs wateroplosbare blaarvoedingsprodukt.

Veggies OEMFF®	g/kg					mg/kg					
	N	P	K	Mg	S	Fe	Mn	Zn	Cu	B	Mo
<b>Starter</b>	147,79	88,86	156,27	13,44	19,42	1607	1339	1236	232	1426	203
<b>Grow</b>	222,03	50,15	128,33	13,46	19,45	1544	1258	1205	231	1370	217
<b>Fruit</b>	81,29	53,36	304,26	13,32	19,24	1611	1299	1221	222	1408	219

## Voordele (Hoekom):

- Die drie **Veggies OEMFF®** produkte is so geformuleer dat dit die drie fenologiese stadia van groentegewasse ondersteun en bevorder. **Veggies OEMFF® Starter** is effektief vroeg in die seisoen waar wortelontwikkeling en die vestiging van die jong plant uiters belangrik is. **Veggies OEMFF® Grow** se elementsamestelling is van so aard dat vegetatiewe groei bevorder word, terwyl **Veggies OEMFF® Fruit** in blom en die selvergrotingsfase van vrugte en knolle 'n belangrike rol speel.
- Alle toepaslike mikro-elemente is gecheleer.

## Gebruike (Waar):

- Veggies OEMFF®** bevat makro-, sekondêre- en mikro-elemente in 'n verhouding wat spesifiek voordelig vir groente is en kan as algemene aanvulling gebruik word.
- Is uiters geskik waar groente aan stremming soos versuip/droogte, koue/hitte of chemiese stremming blootgestel was.
- Kan onder hoë- of lae grond pH toestande gebruik word waar probleme met mikro-elemente verwag word.

## Toediening (Hoe):

Produkt	kg/ha	Tyd van Toediening
<b>Veggies OEMFF® Starter</b>	2kg	Begin 14 dae na opkoms of 7 dae na uitplant. Herhaal 3-4 keer met 10 tot 14 dae intervalle.
<b>Veggies OEMFF® Grow</b>	2kg	Elke 7-14 dae tot begin van blomfase.
<b>Veggies OEMFF® Fruit</b>	2kg	Elke 7-14 dae gedurende blom- en vrugfase.

## Gebruike (Waar):

- Dien **Veggies OEMFF®** toe met 'n hoë volume bespuiting om volledige benutting te verseker. Indien lae volume bespuitings gebruik word, moet 'n maksimum konsentrasie van 2% (2 kg/100 l water) nie oorskry word nie.
- Dien **Veggies OEMFF®** vroeg in die oggend of laat in die middag toe om maksimum opname te verseker.
- Moenie **Veggies OEMFF®** toedien by temperature hoër as 30 °C of indien plante verlep is nie. Dit kan veroorsaak dat opname benadeel word en gewasse beskadig word.
- Dit word sterk aanbeveel dat 'n klein hoeveelheid van die bespuitingsmengsel vooraf gemeng word om verenigbaarheid te evalueer.
- Die gebruik van 'n geskikte kleefmiddel word sterk aanbeveel.

**Let wel: Blaarvoeding moet in kombinasie en as ondersteuning van 'n bemestingsprogram gebruik word.**



« Enhanced efficiency through innovation »



# VEGGIE OEMFF®



## Name:

**Veggie OEMFF® Starter, Veggie OEMFF® Grow, Veggie OEMFF® Fruit**

## Properties (What):

It is a fine, highly soluble foliar product.

Veggie OEMFF®	g/kg					mg/kg					
	N	P	K	Mg	S	Fe	Mn	Zn	Cu	B	Mo
<b>Starter</b>	147,79	88,86	156,27	13,44	19,42	1607	1339	1236	232	1426	203
<b>Grow</b>	222,03	50,15	128,33	13,46	19,45	1544	1258	1205	231	1370	217
<b>Fruit</b>	81,29	53,36	304,26	13,32	19,24	1611	1299	1221	222	1408	219

## Advantages (Why):

- The three **Veggie OEMFF®** products were formulated to support and enhance the three phenological stages of vegetable crops. **Veggie OEMFF® Starter** is effective early in the season where root development and the establishment of the young plant are of the utmost importance. **Veggie OEMFF® Grow** enhances growth because of its specific element composition, while **Veggie OEMFF® Fruit** plays an important role in flowering and cell enlargement of fruit and bulbs.
- All the applicable micro-elements are chelated.

## Uses (Where):

- Veggie OEMFF®** contains macro-, secondary- and micro-elements in a specific ratio which is beneficial to vegetables and can be used as a general application if needed.
- Extremely useful if plants were exposed to stresses like water logged conditions/drought, heat/cold or chemical stress.
- Can be used under high or low pH conditions where micro-nutrient deficiencies are expected.

## Application (How):

Product	kg/ha	Application time
<b>Veggie OEMFF® Starter</b>	2 kg	Start application 14 days after emergence or 7 days after transplant. Repeat 3-4 times with 14 day intervals.
<b>Veggie OEMFF® Grow</b>	2 kg	Apply with 7-14 day intervals until beginning of the flowering stage.
<b>Veggie OEMFF® Fruit</b>	2 kg	Apply with 7-14 day intervals during flowering and fruiting stages.

## Uses (Where):

- Apply **Veggie OEMFF®** as a high volume foliar spray to ensure thorough wetting. When low volume sprayers are used, do not exceed a maximum concentration of 2% (2 kg/100 ℓ water).
- Apply **Veggie OEMFF®** range early in the morning or late afternoon to ensure maximum absorption.
- Do not spray when temperatures exceed 30 °C or onto wilted plants as this can result in reduced uptake and damage to the plants.
- It is advised that an observation test be done by first mixing a small amount of the spray solution.
- The use of a suitable sticker is recommended.

**Note: Foliar sprays should be used in conjunction and support of normal fertilizing practices.**